



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2022

Molecular foundations of prion strain diversity

Carta, Manfredi ; Aguzzi, Adriano

Abstract: Despite being caused by a single protein, prion diseases are strikingly heterogenous. Individual prion variants, known as strains, possess distinct biochemical properties, form aggregates with characteristic morphologies and preferentially seed certain brain regions, causing markedly different disease phenotypes. Strain diversity is determined by protein structure, post-translational modifications and the presence of extracellular matrix components, with single amino acid substitutions or altered protein glycosylation exerting dramatic effects. Here, we review recent advances in the study of prion strains and discuss how a deeper knowledge of the molecular origins of strain heterogeneity is providing a foundation for the development of anti-prion therapeutics.

DOI: <https://doi.org/10.1016/j.conb.2021.07.010>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-206180>

Journal Article

Published Version



The following work is licensed under a Creative Commons: Attribution 4.0 International (CC BY 4.0) License.

Originally published at:

Carta, Manfredi; Aguzzi, Adriano (2022). Molecular foundations of prion strain diversity. *Current Opinion in Neurobiology*, 72:22-31.

DOI: <https://doi.org/10.1016/j.conb.2021.07.010>



Molecular foundations of prion strain diversity

Manfredi Carta and Adriano Aguzzi

Abstract

Despite being caused by a single protein, prion diseases are strikingly heterogeneous. Individual prion variants, known as strains, possess distinct biochemical properties, form aggregates with characteristic morphologies and preferentially seed certain brain regions, causing markedly different disease phenotypes. Strain diversity is determined by protein structure, post-translational modifications and the presence of extracellular matrix components, with single amino acid substitutions or altered protein glycosylation exerting dramatic effects. Here, we review recent advances in the study of prion strains and discuss how a deeper knowledge of the molecular origins of strain heterogeneity is providing a foundation for the development of anti-prion therapeutics.

Addresses

Institute of Neuropathology, University Hospital of Zurich, University of Zurich, Schmelzbergstrasse 12, 8091 Zurich, Switzerland

Corresponding author: Aguzzi, Adriano (adriano.aguzzi@usz.ch)

Current Opinion in Neurobiology 2022, **72**:22–31

This review comes from a themed issue on **Neurobiology of Disease**

Edited by **Bart de Strooper** and **Huda Zoghbi**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online 27 July 2021

<https://doi.org/10.1016/j.conb.2021.07.010>

0959-4388/© 2021 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Introduction

Prion diseases are a group of fatal neurodegenerative disorders caused by prions. These unique infectious agents are not known to contain any nucleic acids and appear to be composed primarily of a misfolded form of the prion protein PrP, which is encoded by the prion protein gene *PRNP* [1,2]. Histologically, prion diseases show vacuolation, astrogliosis, neuroinflammation and deposition of PrP^{Sc}, a misfolded, beta-sheet-rich counterpart of the cellular prion protein PrP^C, a glycosylphosphatidylinositol-anchored (GPI-anchored) protein that is abundantly expressed in the brain. Prions propagate through the templated misfolding of PrP^C to infectious prion aggregates [3].

Most cases of human prion disease are sporadic (sporadic Creutzfeldt–Jakob disease, sCJD), with no known cause. Inherited *PRNP* mutations cause the genetic prion diseases familial Creutzfeldt–Jakob disease, Gerstmann–Sträussler–Scheinker syndrome (GSS) and fatal familial insomnia. Prion diseases acquired via infection include variant CJD and kuru, where prions are acquired via dietary routes, and iatrogenic CJD, caused by accidental inoculation through medical procedures [4].

Although they are caused by a misfolded form of one single protein, prion diseases of animals and humans are strikingly heterogeneous and are associated with a wide range of clinicopathological variants with different symptoms, incubation periods, rates of disease progression, histology, PrP^{Sc} deposit morphologies and anatomical distributions of disease lesions [5–7].

Given that prions are not known to contain nucleic acids, these properties must be encoded in the higher-order aggregate structure. Indeed, prions exist in distinct conformational variants, known as strains, whose properties can be serially transmitted and reproduced in animal models of disease and also in human infection [6,8,9]. Furthermore, human sporadic prion disease strains can also be reproduced by the *in vitro* prion amplification technique protein misfolding cyclic amplification (PMCA), where prions are propagated in the presence of brain homogenate [10].

Individual prion strains produce PrP^{Sc} with distinct biochemical properties, such as the degree of electrophoretic mobility [11], susceptibility to protease digestion, immunoreactivity to antibodies targeting conformational PrP epitopes [12], glycosylation patterns [13] and stability on treatment with chaotropic salts [14] or heat [15].

Prion strains cause pathology with specific anatomical distributions. A study of sCJD with diffusion-weighted magnetic resonance imaging showed that the disease primary site and ordering of lesion propagation were specific to sCJD histotypes, which are caused by distinct strains. For instance, in the most common histotype, MM1, lesions spread from the neocortex to subcortical regions, with the opposite pattern seen in the histotype VV2 (see [Figure 1](#)) [16**]. The observation that many

self-templating prion strains with defined properties exist leads to the question of how their structure was formed in the original prion seed and how the strain-coding information is transmitted during prion replication. Here, we review how recent histological, genetic and biochemical studies are unravelling the molecular origins of prion strain diversity and discuss how these advances are guiding the development of innovative anti-prion therapeutics.

Determinants of strain properties


Prion protein primary structure

The properties of de novo prion seeds are influenced by the primary amino acid sequence of PrP. For instance, a geographically variable polymorphism in codon 129 of *PRNP*, which codes for methionine (M) or valine (V), influences biochemical features of prions when they emerge in the primary host, and these properties can be transmitted with infection [17–19]. This polymorphism also influences the risk of spontaneously developing prion disease [20], age at onset of genetic disease [21,22], clinical and histological features of disease [7] and the incubation period after exposure to prions [23]. In line with the last observation, the overall PrP sequence determines species barriers; while bank voles can be readily infected with strains originating in various

species and bank vole PrP is susceptible to conversion by several human prion strains [24,25], dogs are resistant to prion infection, probably thanks to the presence of a single negatively charged amino acid (aspartic or glutamic acid) in position 163 of canine PrP, a residue not found in other species [26]. Mutations that alter the amino acid sequence of PrP can cause hereditary prion disease. Around 50–60 variants have been described, including missense mutations, truncations [27] and insertions into the octapeptide repeat region of PrP [28,29], a repetitive domain close to the PrP N-terminus likely to be a mediator of prion toxicity [30]. Full penetrance has been established for four missense mutations, which also demonstrably cause the formation of infectious prions, as shown by transmissibility experiments where rodents and non-human primates were inoculated with material obtained from diseased brains of mutation carriers; furthermore, in mice, homologous *Prnp* mutations cause spontaneous disease [31–34]. Individuals with sporadic prion disease are not known to carry any of these pathogenic mutations in the germline.

PRNP polymorphisms can also be protective. A study of persons who were exposed to kuru via cannibalistic funerary practices, but did not develop the disease, led to the discovery of G127V, a resistance factor whose

Figure 1



Prion strains cause different disease phenotypes. PrP^{Sc} aggregate morphology and anatomical spreading patterns (based on the study by Pascuzzo et al. [16**]) are shown in the CJD histotypes MM1 and VV2. MM1 is the main CJD subtype found in individuals homozygous for methionine at *PRNP* codon 129, whereas VV2 is the main type in valine homozygous individuals. Dark colour indicates the 'epicentre' of pathology, with lighter coloured regions affected later in the disease course. (a) Small punctate synaptic PrP^{Sc} deposits in MM1, cerebellar cortex. (b) Plaque-like PrP^{Sc} deposits in VV2, cerebellar cortex. (c) In MM1, diffusion-weighted imaging showed that pathology most often began in the precuneus and parietal lobe, followed by spread to the frontal and cingulate lobes, with the cerebellum and thalamus affected at late stages. (d) In VV2, the cerebellum is usually affected early in the disease, followed by spread to the basal ganglia and thalamus, insular cortex, frontal lobe and parietal lobe, with the occipital cortex affected at a late stage.

emergence was favoured by the selective pressure of the kuru epidemic [35]. Overall, these observations highlight how substituting a single amino acid in PrP can drastically influence the formation and propagation of prion disease, with effects ranging from protective to inexorably fatal. Several genome-wide association studies have attempted to find variants in other genes that influence the risk of developing prion disease, but the rarity of prion disease poses a significant challenge to these efforts. Apart from variants in *PRNP*, the search for associations has been mostly elusive, but weak associations with variants in *STX6* and *GAL3ST1* were reported recently. Risk single nucleotide polymorphisms (SNPs) in *STX6* are also shared with progressive supranuclear palsy, possibly indicative of common causal mechanisms [36**].

Aggregate biophysics

The pathogenicity of prions is associated with certain biophysical properties which lead to the formation of aggregates with characteristic morphologies. Mouse-adapted prions that form smaller, diffuse, subfibrillar aggregates and are unstable under denaturing conditions are particularly pathogenic and more likely to enter the central nervous system through peripheral nerves, whereas plaque-forming prions are likely to be less pathogenic [37]. Indeed, although both subfibrillar and fibrillar strains can be taken up by neurons by macropinocytosis and transported through axons to the cell body, only subfibrillar strains were efficiently transported to the brain. Sonication of fibrillar strains significantly increased solubility and disease attack rate on peripheral inoculation, showing that smaller PrP species are more pathogenic and that promoting fibril stability and assembly could slow prion propagation [38]. This makes sense mechanistically because the prion is defined as the minimal infectious unit, and it logically consists of one single molecular aggregate—whatever its size. Hence, taking a fibril and fragmenting it into a large number of smaller aggregates is necessarily bound to increase the number of infectious particles even if the total amount of prion protein remains unchanged. In humans, small, non-fibrillar prions are also likely to be particularly infectious. Purified protease-resistant GSS aggregates containing PrP amino acids ~90–150 were highly infectious and capable of causing disease on their own, suggesting that this part of the protein is a minimal component required for prion formation [39*].

A potentially protective prion strain has also been described. Anti-prions are self-replicating but innocuous PrP aggregates which can compete with toxic prions for their substrate PrP^C, resulting in its depletion, inhibiting the spread of toxic prions. When given as a prophylactic treatment in hamsters, anti-prions significantly delayed the onset of clinical symptoms after inoculation with a highly pathogenic prion strain. These findings

indicate that prion replication and toxicity are uncoupled and provide a remarkable example of the phenotypic diversity of prion strains [40**].

Cofactors of prion replication

Prions typically form diffuse, synaptic or punctate deposits, whereas the formation of amyloid plaques composed of PrP is relatively rare and typically associated with a longer disease duration; however, certain strains preferentially form plaques when inoculated into wild-type mice [41*]. Therefore, genetic modifications that alter the pattern of prion deposition could be of particular interest. Whether prion strains form diffuse deposits or plaques has been proposed to be influenced by cofactors, particularly by glycosyl residues of PrP, phospholipids and polyanionic compounds such as RNA or extracellular matrix (ECM) components [42]. In particular, glycans stabilise PrP and can inhibit fibril nucleation [43,44].

A recent study explored the effect of the ECM component heparan sulphate (HS) by inoculating prions into mice which produce shortened HS chains. Prolonged survival and redistribution of PrP plaques from brain parenchyma to blood vessels were seen after infection with fibrillar, plaque-forming prions. Plaques consisted of extracellularly assembled, protease-cleaved, GPI-anchorless PrP and were also enriched in HS. Similarly, in sCJD and GSS, plaque-forming prions had the highest amounts of bound HS, whereas lower levels were seen in cases with diffuse deposits. Given that HS deficiency prolonged survival, HS was found to be a disease-modifying cofactor of prion replication that accelerates prion disease progression [45*].

A similar study investigated the significance of PrP glycosylation using 1. subfibrillar prions, 2. plaque-forming prions and 3. prions consisting of GPI-anchored PrP, which were inoculated into mice with different PrP glycosylation profiles. Expression of underglycosylated PrP resulted in prions that are usually subfibrillar forming plaques, which consisted of PrP cleaved by the protease ADAM10 and HS. The opposite effect was seen in mice producing triglycosylated PrP: here, challenge with plaque-forming prions leads to accelerated disease with deposition of diffuse, subfibrillar aggregates. GPI-anchored prions recruited unglycosylated, GPI-anchored PrP and aggregated to large plaques. However, no difference was seen after infection with the third strain consisting of unmodified, GPI-anchored PrP, which formed diffuse, HS-deficient deposits irrespective of altered glycosylation. These findings highlight how glycans, protease cleavage and GPI anchors shape prion disease phenotypes, with a strong influence on aggregate morphology, disease progression and interaction with polyanionic HS being evident. In particular, PrP plaques appeared to be

preferentially formed by underglycosylated PrP, whereas addition of glycosyl residues favoured the deposition of subfibrillar aggregates [41*].

In another study investigating the influence of cofactors on pathogenicity, a protein-only PrP^{Sc} preparation produced from recombinant PrP by the prion-amplification technique PMCA failed to induce disease after inoculation into bank voles, which are highly susceptible to infection by various prion strains. However, if the same protein-only PrP^{Sc} was seeded into a PMCA reaction containing brain homogenate, potentially infectious PrP^{Sc} was produced, indicating that native PrP (not recombinant PrP produced in bacteria) and additional cofactors present within brain homogenate were required for *in vitro* production of highly pathogenic prions [46].

Using PMCA, it was also found that prion strains replicated preferentially in the presence of PrP glycoforms and cofactor molecules that were found in the host species where the prions originated. Given that PrP expression, PrP glycosylation [47], membrane lipids and ECM components vary throughout the brain, strain-specific patterns of neurotropism probably depend on the presence of PrP glycoforms and cofactor molecules in different parts of the brain, that is, prions preferentially propagate in brain regions that provide the substrates required by an individual prion strain for efficient conversion [48*].

Antemortem diagnosis of sCJD is now commonly made with real-time quaking-induced conversion (RT-QuIC) assays, which can detect prions in patients' cerebrospinal fluid with very high sensitivity [49,50]. In RT-QuIC assays, the presence of PrP^{Sc} in the analyte provokes misfolding of PrP^C through cycles of elongation and fragmentation, generating aggregates that can be stained with the fluorescent dye thioflavin T. It is important to understand that the original intuition of breaking aggregates and letting them regrow cyclically, which was implemented in PMCA, predates RT-QuIC by more than a decade. It was Claudio Soto who demonstrated, in a landmark experiment, that this was possible [24]. While RT-QuIC improved technically on the Soto methodology by substituting sonication with quaking, it was an evolutionary refinement rather than a revolutionary invention. As mentioned, PMCA products are demonstrably infectious. To investigate whether RT-QuIC products are pathogenic prions or an innocuous form of aggregated PrP, RT-QuIC products were inoculated into the brains of mice overexpressing human PrP, where they failed to induce clinically apparent disease. However, histological workup of the inoculated brains revealed the presence of abnormal cortical PrP deposits and vacuolation, but no evidence of gliosis was found. RT-QuIC seeding activity was also detected in the inoculated brains. These results indicate that RT-QuIC products can provoke PrP aggregation and limited

cellular pathology *in vivo* but that they represent a species of PrP that is distinct from pathogenic prions [51*]. Nevertheless, the finding that a limited degree of pathology was present is of note and provides further evidence that PrP aggregates encompass a spectrum of many different species, whose degree of pathogenicity ranges from innocuous to highly infectious.

Taken together, these studies of PrP post-translational modification provide compelling evidence for a critical role of PrP glycosylation and of cofactors found in brain homogenate (possibly lipids and ECM components such as HS) in prion propagation, particularly in the formation of infectious aggregates. Furthermore, the spatial distribution of PrP post-translational modifications and of cofactor molecules in the brain might explain why individual strains preferentially seed pathology in certain brain regions.

Structural and biophysical studies

Recent studies have provided unprecedented insight into the molecular structure of prion fibrils. Using cryo-electron microscopy, a denaturant- and protease-resistant prion fibril formed from an unglycosylated fragment of human PrP was shown to contain two intertwined protofilaments with screw symmetry linked by a hydrophobic interface. Based on this structure, certain familial prion disease variants were predicted to form closely related polymorphs and a protective amino acid substitution was predicted to form steric clashes, explaining why it might disrupt prion fibril formation [52*]. A similar structural study in preprint of scrapie prions carrying post-translational modifications demonstrated how glycosyl residues markedly alter fibril structure. Moreover, GPI anchors might cause fibril-membrane interactions that are likely to be highly relevant to prion propagation and pathology [53]. The findings in these structural studies could help explain how prion strain properties are shaped by the PrP amino acid sequence and post-translational modifications, providing a molecular basis for strain diversity.


Novel biophysical and microfluidic diffusional sizing methods have enabled the study of prion propagation kinetics both *in vitro* and *in vivo*. In a recent study, such methods were used to formulate a mathematical framework of prion aggregation kinetics. *In vivo* prion aggregation kinetics of prions were consistent with *in vitro* findings and demonstrated that fragmentation was the dominant mode of prion fibril replication *in vivo* [54*].

Novel approaches to prion genetics

Somatic mutations

The rarity of sporadic prion disease renders it very challenging to use population genetics to uncover novel risk loci. Alternative approaches are therefore needed.

Figure 2



Mechanisms of action of anti-prion therapeutics. RNA coding for PrP is translated, forming PrP^C (blue molecules). At the onset of prion pathology, PrP^C forms a primary prion seed composed of misfolded monomers (red blocks). Aggregates grow by recruiting and misfolding further PrP^C monomers. Prion aggregates fragment (lightning) into smaller daughter prions, which continue to replicate by a cycle of elongation and fragmentation. Prions interact with cell surface PrP^C, initiating toxic signalling pathways. Antisense oligonucleotides bind RNA and cause it to be degraded, inhibiting expression of PrP^C. Anti-prions are non-toxic prion strains which deplete PrP^C, preventing its recruitment into toxic prions. Therapeutic antibodies bind PrP^C, with several beneficial effects: prevention of prions from binding PrP^C, decreased protein misfolding and inhibition of toxic signalling. Therapeutic antibodies and luminescent conjugated polythiophenes (LCPs) stabilise aggregates, preventing fragmentation into daughter prions.

In recent years, several remarkable studies have shown how the human brain is a genomic mosaic, with individual cells harbouring clonal or private mutations that were acquired as a consequence of mutagenic processes, some of them potentially altering the function of affected genes [55,56]. This genomic diversity raises the possibility that somatic mutations in key disease-related genes could contribute to neurodegeneration, including to the pathogenesis of prion disease, and several recent studies have begun to investigate this possibility.

With ageing and in Alzheimer's disease, a number of putatively pathogenic mutations were detected in pathways implicated in hyperphosphorylation of tau, suggesting that they might have contributed to Alzheimer's disease pathology [57*]. A similar study of semantic dementia revealed individuals with putatively pathogenic mosaic mutations in *TARDBP* (with a mutant allele fraction of 1–3%), one of the main genes implicated in the pathogenesis of this neurodegenerative disease [58]. A study of the synucleinopathies Parkinson's disease and multiple system atrophy found that nigral dopaminergic neurons harboured somatic *SNCA* (α -synuclein) copy number gains compared with controls, suggesting that supernumerary *SNCA* copies, possibly leading to overexpression of α -synuclein, might be a significant pathogenetic risk factor [59]. Indeed,

germline locus triplication of *SNCA* is a long-known cause of hereditary Parkinson's disease [60].

Somatic mutations were already proposed to lie at the origin of sporadic prion disease around 30 years ago [61], but there are no reports on large-scale searches for *PRNP* mutations in prion disease samples. Although biosafety hazards and the limited availability of patient samples render it difficult to sequence DNA from the prion-contaminated brain, targeted sequencing of *PRNP* or whole genome sequencing of the somatic genome could potentially uncover known pathogenic variants or novel genetic risk factors.

Screens

Genetic screens might also identify crucial genes hitherto undetected by population genetics. An arrayed whole-transcriptome RNA interference screen in pre-print identified a number of genes regulating PrP^C biosynthesis, with the RNA-binding post-transcriptional repressor Pumilio-1 emerging as a potent regulator causing degradation of *PRNP* mRNA [62]. Similar screens for genes regulating prion propagation or production of potential prion co-factors could help unravel prion pathogenesis and disease heterogeneity. The 'hits' uncovered by such unbiased screens could represent genetic risk loci for sporadic prion disease, potentially containing disease-modifying variants in affected

patients. In a different screen using a short-hairpin RNA library targeting 133 genes related to protein homeostasis, several members of the Hsp70 family of chaperones and related proteins were identified as regulators of PrP^C expression. A molecule inhibiting the Hsp70 family reduced PrP^C levels and also PrP^{Sc} formation [63]. Although this finding could hold promise for the development of small-molecule therapeutics, it might be necessary to find a more specific Hsp70 family member that is of relevance to prion pathophysiology, as the pervasive functions of the Hsp70 family in cell biology might make them too broad a target [64].

Therapies

Next, we discuss how recent findings in the field of prion research could inform the development of anti-prion therapeutics (see the overview in Figure 2).

Aggregate stabilisation

Given that prions replicate via fragmentation, stabilisation of aggregates with molecules that intercalate into the quaternary structure of prions could be a promising therapeutic strategy. Luminescent conjugated polymers bind prions and other amyloids [6,65], stabilise prions [66] and extend the lifespan of prion-infected mice by up to 80%, while being well-tolerated [67]. Given that these molecules are luminescent, imaging approaches could be used to track their function and distribution *in vivo*. Using similar compounds and multiphoton microscopy, longitudinal imaging of the spread of tau and amyloid was achieved *in vivo* [68**]. Similar studies of prions, with longitudinal imaging surveillance of their effects, might help identify promising drug candidates. Polymers that specifically bind prion aggregates could also be conjugated to effector molecules, such as immunological signalling proteins, for specific targeting of aggregates.

Antibodies

Anti-PrP antibodies exert differential effects depending on their targeted epitope. Antibodies binding the C-terminal globular domain of PrP are toxic and reproduce many of the features of prion disease, whereas antibodies binding the N-terminal flexible tail protect against disease, presumably by preventing prions from binding PrP^C and exerting toxic effects [30]. Recent data from our laboratory show that expression of PrP^C in neurons—but not astrocytes—is required for prion toxicity [69]; hence, preventing prions from binding neuronal PrP^C with suitable antibodies might be a viable strategy to prevent neurodegeneration.

Anti-PrP autoantibodies, including clones with therapeutic potential, were recently shown to be found in healthy individuals. No associations with manifest prion disease, *PRNP* mutation status or any other specific pathologies could be identified, indicating that

circulating anti-PrP antibodies are likely to be well-tolerated, making them a viable therapeutic approach [70,71]. Rationally designed therapeutic anti-PrP antibodies have already shown promise in early experimental studies. A bispecific antibody which simultaneously targets both the globular and flexible tail domains of PrP reduced neurotoxicity and the formation of soluble oligomers in a brain slice model of prion disease, possibly by blocking the prion-binding site in the globular domain and stabilising the flexible tail [72]. Further studies, particularly *in vivo* treatments, are required to determine whether anti-PrP antibodies might hold promise as a neuroprotective therapy or possibly also as promoters of prion clearance. In particular, whether limited passage through the blood–brain barrier could be a hindrance to therapy remains to be elucidated.

Antisense oligonucleotides

Gene knockout of mouse *Prnp* produces a demyelinating polyneuropathy phenotype, as PrP is required for myelin maintenance in peripheral nerves [73,74], but in the central nervous system, lack of PrP expression is currently not known to produce significant effects. Consequently, depleting PrP in the brain might be a promising therapeutic strategy to treat prion disease or a prophylactic strategy to prevent *PRNP* mutations from causing outbreak of genetic prion disease (as in the anti-prion approach described previously). A promising strategy to reduce PrP expression is knockdown using antisense oligonucleotides (ASOs), single-stranded DNA fragments that specifically bind their target mRNA, leading to its degradation by RNase H1. Chronic suppression of PrP expression with ASO treatment extended survival of prion-infected mice in both prophylactic and therapeutic settings, even if given shortly before the onset of evident neuropathology [75**].

In a follow-up study, ASO treatment conferred robust dose-dependent benefits in mice infected with 5 different prion strains, with 21% RNA knockdown already extending survival. ASOs even extended survival if given after the onset of clinically evident disease, in which case they attenuated symptoms and lowered concentrations of biomarkers of neuronal injury [76]. Given that ASOs act by depleting the substrate of prion conversion, their action is independent of strain properties, meaning that they could be effective in all forms of prion disease. This is in contrast to stabilising agents such as luminescent conjugated polymers, which would probably only be effective against certain strains. Furthermore, ASOs could potentially be designed to specifically knock down mutant variants of *PRNP* and be used as a prophylactic treatment of *PRNP* mutation carriers, which might delay or even prevent the outbreak of disease. Combination therapies with ASOs and stabilising agents or antibodies might be

particularly promising. Stabilising agents would confer the additional benefit of inhibiting prion fragmentation, whereas antibodies might enhance prion clearance or prevent the toxic cascade initiated by prions on binding of PrP^C. Overall, combination therapies might give microglia time to clear prions, significantly prolonging survival.

Conclusion

The observation that prion strains composed of a single protein are capable of causing a diverse range of pathological phenotypes has baffled scientists for many decades. Recent advances have brought unprecedented insight into the genetic and biochemical phenomena that shape prion strain diversity. Even though prions are believed to principally consist of protein, the role of non-protein cofactors such as glycans, polyanionic macromolecules and GPI anchors in prion propagation is becoming increasingly apparent. Furthermore, structural studies are elucidating the influence of mutations and post-translational modifications on the molecular architecture of prions. Although prion propagation is increasingly understood—with aggregate fragmentation being shown to play a critical role—how prions begin their existence in the first place is still shrouded in mystery. Here, emerging fields such as the study of genomic heterogeneity and high-content genetic screens might provide further insight into the origin and propagation of prions. A deeper understanding of prion diversity is also likely to inform design of specific anti-prion therapeutics, with approaches that stabilise aggregates, target critical prion epitopes or deplete prion protein holding particular promise.

Conflict of interest statement

Nothing declared.

Acknowledgements

The authors thank Dr Karl Frontzek for assistance with histological imaging. M.C. is supported by the Swiss National Science Foundation. A.A. is supported by the European Research Council, the Swiss National Science Foundation, the Gelu Foundation, the Nomis Foundation, the Swiss Personalized Health Network, the USZ Foundation and the estate of Dr. Hans Salvisberg.

References

Papers of particular interest, published within the period of review, have been highlighted as:

* of special interest

** of outstanding interest

- Bolton David C, McKinley Michael P, Prusiner Stanley B: **Identification of a protein that purifies with the scrapie prion.** *Science* 1982, **218**. ISSN: 0036-8075:1309–1311.
- Basler K, Oesch B, Scott M, Westaway D, Wälchli M, Groth DF, McKinley MP, Prusiner SB, Weissmann C: **Scrapie and cellular PrP isoforms are encoded by the same chromosomal gene.** *Cell* 1986, **46**. ISSN: 0092-8674:417–428.
- Schechel C, Aguzzi A: **Prions, prionoids and protein misfolding disorders.** *Nat Rev Genet* 2018, **19**. ISSN: 1471-0056: 405–418, <https://doi.org/10.1038/s41576-018-0011-4>.
- Aguzzi A, Calella AM: **Prions: protein aggregation and infectious diseases.** *Physiol Rev* 2009, **89**. ISSN: 0031-9333: 1105–1152, <https://doi.org/10.1152/physrev.00006.2009> (Print) 0031-9333.
- Collis SC, Kimberlin RH: **Long-term persistence of scrapie infection in mouse spleens in the absence of clinical disease.** *FEMS (Fed Eur Microbiol Soc) Microbiol Lett* 1985, **29**:111–114.
- Sigurdson Christina J, Nilsson KPeter R, Hornemann Simone, Manco Giuseppe, Polymenidou Magdalini, Schwarz Petra, Leclerc Mario, Hammarström Per, Kurt Wüthrich, Aguzzi Adriano: **Prion strain discrimination using luminescent conjugated polymers.** *Nat Methods* 2007, **4**:1023–1030.
- Parchi Piero, De Boni Laura, Saverioni Daniela, Cohen Mark L, Ferrer Isidro, Gambetti Pierluigi, Gelpi Ellen, Giaccone Giorgio, Jean-Jacques Hauw, Höftberger Romana, et al.: **Consensus classification of human prion disease histotypes allows reliable identification of molecular subtypes: an inter-rater study among surveillance centres in Europe and USA.** *Acta Neuropathol* 2012, **124**:517–529.
- Hill Andrew F, Desbruslais Melanie, Joiner Susan, Katie CL Sidle, Gowland Ian, Collinge John, J Doey Lawrence, Lantos Peter: **The same prion strain causes vCJD and BSE.** *Nature* 1997, **389**: 448–450.
- Moda Fabio, Suardi Silvia, Giuseppe Di Fede, Indaco Antonio, Limido Lucia, Vimercati Chiara, Ruggerone Margherita, Campagnani Ilaria, Jan Langeveld, Terruzzi Alessandro, et al.: **MM2-Thalamic Creutzfeldt–Jakob disease: neuropathological, biochemical and transmission studies identify a distinctive prion strain.** *Brain Pathol* 2012, **22**:662–669.
- Bélondrade Maxime, Simon Nicot, Mayran Charly, Bruyere-Ostells Lilian, Almela Florian, Michele A Di Bari, Levavasseur Etienne, Watts Joel C, Fournier-Wirth Chantal, Lehmann Sylvain, et al.: **Sensitive protein misfolding cyclic amplification of sporadic Creutzfeldt–Jakob disease prions is strongly seed and substrate dependent.** *Sci Rep* 2021, **11**: 1–12.
- Parchi Piero, Giese Armin, Capellari Sabina, Brown Paul, Schulz-Schaeffer Walter, Otto Windl, Zerr Inga, Herbert Budka, Kopp Nicolas, Pedro Piccardo, et al.: **Classification of sporadic Creutzfeldt–Jakob disease based on molecular and phenotypic analysis of 300 subjects.** *Ann Neurol* 1999, **46**:224–233.
- Polymenidou Magdalini, Stoeck Katharina, Glatzel Markus, Martin Vey, Bellon Anne, Aguzzi Adriano: **Coexistence of multiple PrP^{Sc} types in individuals with Creutzfeldt–Jakob disease.** *Lancet Neurol* 2005, **4**:805–814.
- Hill Andrew F, Joiner Susan, Beck Jonathan A, Campbell Tracy A, Dickinson Andrew, Poulter Mark, Wadsworth Jonathan DF, Collinge John: **Distinct glycoform ratios of protease resistant prion protein associated with PRNP point mutations.** *Brain* 2006, **129**:676–685.
- Pirsinu Laura, Di Bari Michele, Marcon Stefano, Vaccari Gabriele, D'Agostino Claudia, Fazzi Paola, Esposito Elena, Galeno Roberta, Langeveld Jan, Agrimi Umberto, et al.: **A new method for the characterization of strain-specific conformational stability of protease-sensitive and protease-resistant PrP^{Sc}.** *PLoS One* 2010, **5**, e12723.
- Marín-Moreno Alba, Aguilar-Calvo Patricia, Moudjou Mohammed, Espinosa Juan Carlos, Béringue Vincent, Torres Juan María: **Thermostability as a highly dependent prion strain feature.** *Sci Rep* 2019, **9**:1–10.
- Pascuzzo Riccardo, Oxtoby Neil P, Young Alexandra L, Blevins Janis, Castelli Gianmarco, Garbarino Sara, Cohen Mark L, Schonberger Lawrence B, Gambetti Pierluigi, Appleby Brian S, et al.: **Prion propagation estimated from brain diffusion MRI is subtype dependent in sporadic Creutzfeldt–Jakob disease.** *Acta Neuropathol* 2020, **140**:169–181.

In diffusion-weighted imaging (DWI) MRI of sCJD patients, lesion primary site and spreading pattern was shown to correlate with sCJD histotype. This reproducible spreading pattern might indicate that prion neurotropism depends on the presence of anatomically variable factors which specifically favour propagation of certain strains.

- Soldevila Marta, Calafell Francesc, Andrés Aida M, Yagüe Jordi, Helgason Agnar, Stefánsson Kári, Bertranpetit Jaume: **Prion**

- susceptibility and protective alleles exhibit marked geographic differences.** *Hum Mutat* 2003, **22**:104–105.
18. Tahiri-Alaoui Abdessamad, Gill Andrew C, Disterer Petra, James William: **Methionine 129 variant of human prion protein oligomerizes more rapidly than the valine 129 variant: implications for disease susceptibility to Creutzfeldt-Jakob disease.** *J Biol Chem* 2004, **279**:31390–31397.
 19. Baskakov Ilia, Disterer Petra, Breydo Leonid, Shaw Michael, Gill Andrew, James William, Tahiri-Alaoui Abdessamad: **The presence of valine at residue 129 in human prion protein accelerates amyloid formation.** *FEBS Lett* 2005, **579**: 2589–2596.
 20. Palmer Mark S, J Dryden Aidan, Hughes J Trevor, Collinge John: **Homozygous prion protein genotype predisposes to sporadic Creutzfeldt-Jakob disease.** *Nature* 1991, **352**. ISSN: 1476-4687:340–342.
 21. Mead S, Webb TEF, Campbell TA, Beck J, Linehan JM, Rutherford S, Joiner S, Wadsworth JDF, Heckmann J, Wroe S: **Inherited prion disease with 5-OPRI: phenotype modification by repeat length and codon 129.** *Neurology* 2007, **69**. ISSN: 0028-3878:730–738.
 22. Webb TEF, Poulter M, Beck J, Uphill J, Adamson G, Campbell T, Linehan J, Powell C, Brandner S, Pal S: **Phenotypic heterogeneity and genetic modification of P102L inherited prion disease in an international series.** *Brain* 2008, **131**. ISSN: 0006-8950:2632–2646.
 23. Peter Rudge, Jaunmuktane Zane, Adlard Peter, Bjurström Nina, Caine Diana, Lowe Jessica, Norsworthy Penny, Hummerich Holger, Druey Ron, Wadsworth Jonathan DF: **Latrogenic CJD due to pituitary-derived growth hormone with genetically determined incubation times of up to 40 years.** *Brain* 2015, **138**. ISSN: 1460-2156:3386–3399.
 24. Saborio Gabriela P, Bruno Permann, Soto Claudio: **Sensitive detection of pathological prion protein by cyclic amplification of protein misfolding.** *Nature* 2001, **411**:810–813.
 25. Nonno Romolo, Notari Silvio, Angelo Di Bari Michele, Cali Ignazio, Pirisinu Laura, d'Agostino Claudia, Cracco Laura, Kofsky Diane, Vanni Ilaria, Lavrich Jody, et al.: **Variable protease-sensitive prionopathy transmission to bank voles.** *Emerg Infect Dis* 2019, **25**:73.
 26. Vidal Enric, Fernández-Borges Natalia, Eraña Hasier, Parra Beatriz, Pintado Belén, Sánchez-Martín Manuel A, Charco Jorge M, Ordóñez Montserrat, Pérez-Castro Miguel A, Pumarola Martí, et al.: **Dogs are resistant to prion infection, due to the presence of aspartic or glutamic acid at position 163 of their prion protein.** *FASEB J* 2020, **34**: 3969–3982.
 27. Choi Jin-Kyu, Cali Ignazio, Surewicz Krystyna, Kong Qingzhong, Gambetti Pierluigi, Surewicz Witold K: **Amyloid fibrils from the N-terminal prion protein fragment are infectious.** *Proc Natl Acad Sci USA* 2016, **113**:13851–13856.
 28. Kaski Diego N, Pennington Catherine, Beck Jon, Poulter Mark, Uphill James, Bishop Matthew T, Linehan Jacqueline M, O'Malley Catherine, Wadsworth Jonathan DF, Joiner Susan: **Inherited prion disease with 4-octapeptide repeat insertion: disease requires the interaction of multiple genetic risk factors.** *Brain* 2011, **134**. ISSN: 1460-2156:1829–1838.
 29. Sánchez-Valle Raquel, Yagüe Jordi, Turón Antoni, I Aróstegui Juan, Nos Carlos, Rey MaJesús, Ferrer Isidro, Gelpi Ellen: **Inherited prion disease with 4-octapeptide repeat insertion linked to valine at codon 129.** *Brain* 2012, **135**. ISSN: 1460-2156:e212.
 30. Sonati Tiziana, Reimann Regina R, Falsig Jeppe, Kumar Baral Pravas, O'Connor Tracy, Hornemann Simone, Yaganoglu Sine, Li Bei, Herrmann Uli S, Wieland Barbara: **The toxicity of antiprion antibodies is mediated by the flexible tail of the prion protein.** *Nature* 2013, **501**. ISSN: 1476-4687: 102–106.
 31. Mastrianni JA, Capellari S, Telling GC, Han D, Bosque P, Prusiner SB, DeArmond SJ: **Inherited prion disease caused by the V210I mutation: transmission to transgenic mice.** *Neurology* 2001, **57**. ISSN: 0028-3878:2198–2205.
 32. Tateishi J, Kitamoto T, Hoque MZ, Furukawa H: **Experimental transmission of Creutzfeldt-Jakob disease and related diseases to rodents.** *Neurology* 1996, **46**. ISSN: 0028-3878: 532–537.
 33. Brown Paul, Gibbs Jr CJ, Rodgers-Johnson Pamela, Asher David M, Sulima Michael P, Bacote Alfred, Goldfarb Lev G, Gajdusek D Carleton: **Human spongiform encephalopathy: the National Institutes of Health series of 300 cases of experimentally transmitted disease.** *Ann Neurol: Off J Am Neurol Assoc Child Neurol Soc* 1994, **35**. ISSN: 0364-5134:513–529.
 34. Minikel EV, et al.: **Quantifying prion disease penetrance using large population control cohorts.** *Sci Transl Med* 2016, **8**. ISSN: 1946-6234, <https://doi.org/10.1126/scitranslmed.aad5169>. 322ra9.
 35. Mead Simon, Whitfield Jerome, Poulter Mark, Shah Pares, Uphill James, Campbell Tracy, Al-Dujaily Huda, Hummerich Holger, Beck Jon, Mein Charles A: **A novel protective prion protein variant that colocalizes with kuru exposure.** *N Engl J Med* 2009, **361**. ISSN: 0028-4793:2056–2065.
 36. Jones Emma, Hummerich Holger, Viré Emmanuelle, Uphill James, Dimitriadis Athanasios, Helen Speedy, Campbell Tracy, Norsworthy Penny, Quinn Liam, Whitfield Jerome, et al.: **Genome-wide association study identifies risk variants for sporadic Creutzfeldt-Jakob disease in STX6 and GAL3ST1.** *medRxiv* 2020.
- While several genetic studies failed to show an association between sporadic prion disease risk and genomic loci other than *PRNP*, this recent GWAS found weak but statistically robust associations implicating intracellular trafficking and sphingolipid metabolism.
37. Bett Cyrus, Joshi-Barr Shivanjali, Lucero Melanie, Trejo Margarita, Liberski Pawel, Kelly Jeffery W, Masliah Eliezer, Christina J Sigurdson: **Biochemical properties of highly neuroinvasive prion strains.** *PLoS Pathog* 2012, **8**, e1002522.
 38. Bett Cyrus, Lawrence Jessica, Kurt Timothy D, Orru Christina, Aguilar-Calvo Patricia, Kincaid Anthony E, Witold K Surewicz, Caughey Byron, Wu Chengbiao, Sigurdson Christina J: **Enhanced neuroinvasion by smaller, soluble prions.** *Acta Neuropathol Commun* 2017, **5**:1–11.
 39. Vanni Ilaria, Pirisinu Laura, Acevedo-Morantes Claudia, Kamali-Jamil Razieh, Rathod Vineet, Angelo Di Bari Michele, D'Agostino Claudia, Marcon Stefano, Esposito Elena, Riccardi Geraldina, et al.: **Isolation of infectious, non-fibrillar and oligomeric prions from a genetic prion disease.** *Brain* 2020, **143**:1512–1524.
- The unusually small protease-resistant PrP^{Sc} core found in a certain type of GSS, which forms non-fibrillar particles, was found to be highly infectious, suggesting that it represents the minimal component of PrP that is required for prion infectivity.
40. Diaz-Espinoza Rodrigo, Morales Rodrigo, Concha-Marambio Luis, Moreno-Gonzalez Ines, Moda Fabio, Soto Claudio: **Treatment with a non-toxic, self-replicating anti-prion delays or prevents prion disease in vivo.** *Mol Psychiatry* 2018, **23**:777–788.
- The authors created an "anti-prion", an unusual form of PrP^{Sc} which was self-replicating but innocuous. Anti-prions competed with pathogenic prions for PrP, leading to a striking reduction in infectivity. Anti-prions could represent a potential therapeutic strategy which depletes the substrate of pathogenic prions.
41. Sevillano Alejandro M, Aguilar-Calvo Patricia, Kurt Timothy D, Lawrence Jessica A, Soldau Katrin, Nam Thu H, Schumann Taylor, Pizzo Donald P, Nyström Sofie, Choudhury Biswa, et al.: **Prion protein glycans reduce intracerebral fibril formation and spongiosis in prion disease.** *J Clin Invest* 2020, **130**.
- Mice expressing PrP with altered glycosylation profiles showed markedly different pathology compared to wildtype mice, highlighting how post-translational modifications and polyanionic co-factors shape prion disease phenotype and strain diversity.
42. Caughey Byron, Baron Gerald S: **Prions and their partners in crime.** *Nature* 2006, **443**:803–810.
 43. Wu Emilia L, Qi Yifei, Park Soohyung, Mallajosyula Sairam S, MacKerell Jr Alexander D, Klauda Jeffery B, Im Wonpil: **Insight into early-stage unfolding of GPI-anchored human prion protein.** *Biophys J* 2015, **109**:2090–2100.

44. Mishra Rajesh, Elgland Mathias, Begum Afshan, Fyrner Timmy, Peter Konradsson, Nyström Sofie, Hammarström Per: **Impact of N-glycosylation site variants during human PrP aggregation and fibril nucleation.** *Biochim Biophys Acta - Protein Proteomics* 2019, **1867**:909–921.
 45. Aguilar-Calvo Patricia, Sevillano Alejandro M, Bapat Jaidev, Soldau Katrin, Sandoval Daniel R, Altmeyen Hermann C, Linsenmeier Luise, Pizzo Donald P, Geschwind Michael D, Sanchez Henry, *et al.*: **Shortening heparan sulfate chains prolongs survival and reduces parenchymal plaques in prion disease caused by mobile, ADAM10-cleaved prions.** *Acta Neuropathol* 2020, **139**:527–546.
- Mice which produce short heparan sulfate chains show prolonged survival upon prion infection. Prion plaques were composed of anchorless, cleaved prion protein and showed an altered deposition pattern, which was associated with increased heparan sulfate binding to prion fibrils.
46. Burke Cassandra M, Walsh Daniel J, Steele Alexander D, Agrimi Umberto, Angelo Di Bari Michele, Watts Joel C, Supattapone Surachai: **Full restoration of specific infectivity and strain properties from pure mammalian prion protein.** *PLoS Pathog* 2019, **15**, e1007662.
 47. Makarava Natalia, Chang Jennifer Chen-Yu, Baskakov Ilia V: **Region-specific sialylation pattern of prion strains provides novel insight into prion neurotropism.** *Int J Mol Sci* 2020, **21**: 828.
 48. Burke Cassandra M, Walsh Daniel J, Mark Kenneth MK, Deleault Nathan R, Nishina Koren A, Agrimi Umberto, Di Bari Michele A, Supattapone Surachai: **Cofactor and glycosylation preferences for in vitro prion conversion are predominantly determined by strain conformation.** *PLoS Pathog* 2020, **16**, e1008495.
- This study reports that prion strains replicate preferentially in the presence of PrP glycoforms that were found in the host species where the prions originated, suggesting that prion neurotropism is influenced by differential glycosylation patterns and possibly also by the anatomical distribution of other co-factors.
49. Atarashi Ryuichiro, Satoh Katsuya, Sano Kazunori, Fuse Takayuki, Yamaguchi Naohiro, Ishibashi Daisuke, Matsubara Takehiro, Nakagaki Takehiro, Yamanaka Hitoki, Shirabe Susumu, *et al.*: **Ultrasensitive human prion detection in cerebrospinal fluid by real-time quaking-induced conversion.** *Nat Med* 2011, **17**:175.
 50. Franceschini Alessia, Baiardi Simone, Hughson Andrew G, McKenzie Neil, Moda Fabio, Rossi Marcello, Capellari Sabina, Green Allison, Giaccone Giorgio, Caughey Byron, *et al.*: **High diagnostic value of second generation CSF RT-QuIC across the wide spectrum of CJD prions.** *Sci Rep* 2017, **7**:1–8.
 51. Raymond Gregory J, Race Brent, Orr Christina D, Raymond Lynne D, Bongianini Matilde, Fiorini Michele, Groveman Bradley R, Ferrari Sergio, Sacchetto Luca, Hughson Andrew G, *et al.*: **Transmission of CJD from nasal brushings but not spinal fluid or RT-QuIC product.** *Ann Clin Transl Neurol* 2020, **7**:932–944.
- Real-time quaking induced conversion (RT-QuIC) is a highly sensitive prion detection assay that relies on the ability of PrP^{Sc} to seed aggregates *in vitro*. This study shows that while RT-QuIC products can also seed PrP aggregation *in vivo* upon inoculation into transgenic mice, they are distinct from pathogenic prions and are unlikely to be similarly biohazardous.
52. Glynn Calina, Sawaya Michael R, Ge Peng, Gallagher-Jones Marcus, Short Connor W, Bowman Ronquiahah, Apostol Marcin, Zhou Z Hong, Eisenberg David S, Rodriguez Jose A: **Cryo-EM structure of a human prion fibril with a hydrophobic, protease-resistant core.** *Nat Struct Mol Biol* 2020, **27**:417–423.
- The cryo-EM structure of a denaturant- and protease-resistant prion fibril is presented, permitting predictive studies of the structural effects of PRNP polymorphisms.
53. Kraus Allison, Hoyt Forrest, Schwartz Cindi, Hansen Bryan, Hughson Andrew G, Artikis Efrosini, Race Brent, Caughey Byron: **Structure of an infectious mammalian prion.** *bioRxiv* 2021.
 54. Meisl Georg, Kurt Timothy, Condado-Morales Itzel, Bett Cyrus, Sorce Silvia, Nuvolone Mario, *et al.*: **Scaling analysis reveals the mechanism and rates of prion replication in vivo.** In *Nature Structural & Molecular Biology* 28.4, pp. 365–372; 2021.
- The first study of prion propagation kinetics *in vivo*.
55. Lodato Michael A, Woodworth Mollie B, Lee Semin, D Evrony Gilad, Mehta Bhaven K, Amir Karger, Lee Soohyun, Chittenden Thomas W, Alissa M D'Gama, Cai Xuyu: **Somatic mutation in single human neurons tracks developmental and transcriptional history.** *Science* 2015, **350**. ISSN: 0036-8075: 94–98.
 56. Rodin Rachel E, Dou Yanmei, Kwon Minseok, Sherman Maxwell A, Alissa M D'Gama, Doan Ryan N, Rento Lariza M, Kelly M Girsakis, Bohrsen Craig L, Kim Sonia N, *et al.*: **The landscape of somatic mutation in cerebral cortex of autistic and neurotypical individuals revealed by ultra-deep whole-genome sequencing.** *Nat Neurosci* 2021, **24**:176–185.
 57. Park Jun Sung, Lee Junehawk, Sun Jung Eun, Kim Myeong-Heui, Kim Il Bin, Son Hyeonju, Kim Sangwoo, Kim Sanghyeon, Young Mok Park, Mook-Jung Inhee: **Brain somatic mutations observed in Alzheimer's disease associated with aging and dysregulation of tau phosphorylation.** *Nat Commun* 2019, **10**. ISSN: 2041-1723:1–12.
- Whole exome sequencing of brain tissue revealed that somatic mutations were enriched with age and in Alzheimer's disease patients. Some mutations were found in genes implicated in tau hyperphosphorylation, suggesting a pathophysiological link.
58. van Rooij Jeroen, Mol Merel O, Melhem Shamiram, van der Wal Pelle, Arp Pascal, Paron Francesca, Kaat Laura Donker, Seelaar Harro, Bank Netherlands Brain, Miedema Suzanne SM, *et al.*: **Somatic TARDBP variants as a cause of semantic dementia.** *Brain* 2020, **143**:3827–3841.
 59. Mokretar Katya, Pease Daniel, Taanman Jan-Willem, Soenmez Aynur, Ejaz Ayesha, Lashley Tammaryn, Ling Helen, Gentleman Steve, Houlden Henry, Holton Janice L, *et al.*: **Somatic copy number gains of α -synuclein (SNCA) in Parkinson's disease and multiple system atrophy brains.** *Brain* 2018, **141**:2419–2431.
 60. Singleton AB, Farrer M, Johnson J, Singleton A, Hague S, Kachergus J, Hulihan M, Peuralinna T, Dutra A Nussbaum, Lincoln S, *et al.*: **α -Synuclein locus triplication causes Parkinson's disease.** *Science* 2003, **302**:841–842.
 61. Prusiner Stanley B: **Scrapie prions.** *Annu Rev Microbiol* 1989, **43**:345–374.
 62. Heinzer Daniel, Avar Merve, Pease Daniel Patrick, Dhingra Ashutosh, Yin Jiang-An, Schaper Elke, Dogancay Berre, Emmenegger Marc, Anna Spinelli, Maggi Kevin, *et al.*: **The landscape of PrPC biosynthesis revealed by an arrayed genome-wide interference screen.** *bioRxiv* 2021.
 63. Abrams Jennifer, Taylor Arhar, Mok Sue Ann, Taylor Isabelle R, Kampmann Martin, Gestwicki Jason E: **Functional genomics screen identifies proteostasis targets that modulate prion protein (PrP) stability.** *Cell Stress and Chaperones* 2021, **26**: 443–452.
 64. Rosenzweig Rina, Nillegoda Nadinath B, Mayer Matthias P, Bukau Bernd: **The Hsp70 chaperone network.** *Nat Rev Mol Cell Biol* 2019, **20**:665–680.
 65. Nilsson KPeter R, Ikenberg Kristian, Åslund Andreas, Fransson Sophia, Konradsson Peter, Röcken Christoph, Moch Holger, Aguzzi Adriano: **Structural typing of systemic amyloidosis by luminescent-conjugated polymer spectroscopy.** *Am J Pathol* 2010, **176**:563–574.
 66. Margalith Ilan, Suter Carlo, Ballmer Boris, Schwarz Petra, Tiberi Cinzia, Sonati Tiziana, Falsig Jeppe, Nyström Sofie, Hammarström Per, Åslund Andreas, *et al.*: **Polythiophenes inhibit prion propagation by stabilizing prion protein (PrP) aggregates.** *J Biol Chem* 2012, **287**:18872–18887.
 67. Herrmann Uli S, Anne K Schütz, Hamid Shirani, Huang Danzhi, Saban Dino, Nuvolone Mario, Li Bei, Ballmer Boris, Åslund Andreas Ko, Mason Jeffrey J: **Structure-based drug design identifies polythiophenes as antiprion compounds.** *Sci Transl Med* 2015, **7**. ISSN: 1946-6234. 299ra123–299ra123.
 68. Calvo-Rodriguez Maria, Hou Steven S, Snyder Austin C, Dujardin Simon, Hamid Shirani, Nilsson KPeter R, Bacskai Brian

J: **In vivo detection of tau fibrils and amyloid β aggregates with luminescent conjugated oligothiophenes and multiphoton microscopy.** *Acta Neuropathol Commun* 2019, **7**:1–10.

This paper demonstrates *in vivo* longitudinal imaging of tau fibrils and amyloid β aggregates using luminescent conjugated oligothiophenes and multiphoton microscopy.

69. Asvin KK Lakkaraju, Sorce Silvia, Senatore Assunta, Nuvolone Mario, Guo Jingjing, Schwarz Petra, Moos Rita, Pelczar Pawel, Aguzzi Adriano: **Glial activation in prion diseases is strictly nonautonomous and requires neuronal PrPSc.** *bioRxiv* 2021.
70. Frontzek Karl, Carta Manfredi, Losa Marco, Epskamp Mirka, Meisl Georg, Anane Alice, Brandel Jean-Philippe, Camenisch Ulrike, Castilla Joaquín, Haik Stéphane, *et al.*: **Autoantibodies against the prion protein in individuals with PRNP mutations.** *Neurology* 2020, **95**:e2028–e2037.
71. Senatore Assunta, Frontzek Karl, Emmenegger Marc, Chincisan Andra, Losa Marco, Reimann Regina, Horny Geraldine, Guo Jingjing, Fels Sylvie, Sorce Silvia, *et al.*: **Protective anti-prion antibodies in human immunoglobulin repertoires.** *EMBO Mol Med* 2020, **12**, e12739.
72. Bardelli Marco, Frontzek Karl, Simonelli Luca, Hornemann Simone, Pedotti Mattia, Federica Mazzola, Carta Manfredi, Eckhardt Valeria, D'Antuono Rocco, Virgilio Tommaso, *et al.*: **A bispecific immunotweezer prevents soluble PrP oligomers and abolishes prion toxicity.** *PLoS Pathog* 2018, **14**, e1007335.
73. Bremer Juliane, Baumann Frank, Tiberi Cinzia, Wessig Carsten, Fischer Heike, Schwarz Petra, Steele Andrew D, Toyka Klaus V, Nave Klaus-Armin, Weis Joachim, *et al.*: **Axonal prion protein is required for peripheral myelin maintenance.** *Nat Neurosci* 2010, **13**:310–318.
74. Küffer Alexander, Lakkaraju Asvin KK, Mogha Amit, Petersen Sarah C, Airich Kristina, Doucerain Cédric, Marpakwar Rajlakshmi, Bakirci Pamela, Senatore Assunta, Monnard Arnaud, *et al.*: **The prion protein is an agonistic ligand of the G protein-coupled receptor Adgrg6.** *Nature* 2016, **536**:464–468.
75. Raymond Gregory J, Zhao Hien Tran, Race Brent, Raymond Lynne D, Williams Katie, E Swayze Eric, Graffam Samantha, Le Jason, Caron Tyler, Jacquelyn Stathopoulos, *et al.*: **Antisense oligonucleotides extend survival of prion-infected mice.** *JCI Insights* 2019, **4**.

Targeting PrP mRNA with antisense oligonucleotides (ASOs) lowered PrP expression and extended survival of prion-infected mice.

76. Minikel Eric Vallabh, Zhao Hien T, Le Jason, O'Moore Jill, Pitstick Rose, Graffam Samantha, Carlson George A, Kavanaugh Michael P, Kriz Jasna, Kim Jae Beom, *et al.*: **Prion protein lowering is a disease-modifying therapy across prion disease stages, strains and endpoints.** *Nucleic Acids Res* 2020, **48**:10615–10631.